What if...

## you get bleeding on or



## around your stoma?

## What is bleeding on or around your stoma?

When your stoma was formed, a piece of bowel was brought out onto the surface of your tummy through your abdominal wall. This piece of bowel was then folded back upon itself to form your stoma, rather like when you turn over the cuff of your jumper, so the inner surface or lining of your bowel is now on the outside. This inner surface has lots of superficial blood vessels which can bleed easily, because of this

# What can cause bleeding on or around your stoma?

- Trauma, often the result of over diligent cleaning techniques
- **Friction**
- Clips or stitches around your stoma
- Ulcers
- Granulomas\*
- A ruptured blood vessel...

\*Granulomas are warty growths around the base of your stoma. They probably occur as a reaction to the continual friction from your stoma pouch. Granulomas are harmless but can be tender. Their persistent bleeding can sometimes hinder the application and adhesion of your pouch.



## What if...

## you get bleeding on or

### around your stoma?

#### **Hints and tips**

- Remember, gently does it. Soft cleaning wipes and warm water are all you need to clean around the stoma
- The use of OstoZYME odour neutralising pouch lubricant on the inside of your pouch can help prevent friction
- Sometimes ulcers on your stoma can occur when you have been poorly or stressed and will usually clear up as your health improves
- Make sure that you have at least a 1 mm clearance between your stoma and the pouch to prevent friction occurring
- You may wish to apply a thin layer of stoma paste, a stoma seal or a collar around the base of your stoma for added protection
- Apply cold pressure. A small bag of ice is ideal if you have a specific bleeding point on your stoma. Seek urgent medical advice if this does not stop the bleeding.

